

Symposium  
Brief Alcohol Interventions throughout the health care system

# Brief Alcohol Intervention For Hazardous Drinkers Admitted to the Emergency Department: A randomized controlled trial

Dr Jean-Bernard Daepfen  
Alcohol Treatment Center  
Lausanne University Medical School,  
Lausanne, Switzerland.

ISBRA 2006



centre de traitement en alcoologie

# Research Team

- Co-investigators :
  - ◆ Jacques Gaume, MA
  - ◆ Pierre Bady, PhD
  - ◆ Gerhard Gmel, PhD
  - ◆ Pr. Bertrand Yersin, MD
  - ◆ Pr. Jean-Claude Givel, MD
  - ◆ Dr Jean-Marie Calmes, MD
- Research staff :
  - ◆ Sascha Asal, MA
  - ◆ Muriel Blanc, MD
  - ◆ Nicole Fafard, RN
  - ◆ Muriel Gaulis, MA
  - ◆ Silvia Gibellini, MA
  - ◆ Nicole Michelet, MA
  - ◆ Yves Montagrin, MA
  - ◆ Flavia Prioni-Koetgen, MA
  - ◆ Alicia Seneviratne, MA



# Background

- Brief alcohol intervention (BAI) reduces hazardous drinking in various medical settings, particularly in primary care (Saitz et al, 2006; Bertholet et al, 2004).
- Emergency department (ED) admission offers an opportunity to conduct BAI, but its efficacy in this setting is controversial.

# Published BAI studies involving ED patients

Several studies suggest some efficacy of BAI with ED patients **on drinking outcomes**

- ◆ Chafetz et al, 1962
- ◆ Bernstein et al, 1997
- ◆ Wright et al, 1998
- ◆ Anti-Poiuka et al, 1998
- ◆ Gentilelo et al, 1999
- ◆ Monti et al, 1999
- ◆ Smith et al, 2003
- ◆ Spirito et al, 2004
- ◆ Bazargan-Hejazi et al, 2005

Additional studies suggest some efficacy of BAI with ED patients **on alcohol-related outcomes** (reduction alcohol-related accidents...)

- ◆ Bernstein et al, 1997
- ◆ Monti et al, 1999
- ◆ Forsberg et al, 2000
- ◆ Johnston et al, 2002
- ◆ Nordquist et al, 2005

Only **5 randomized controlled** BAI studies evaluated the efficacy of BAI in ED, 2 of them found beneficial effects on drinking outcomes

- ◆ **Smith et al, 2003**
- ◆ **Spirito et al, 2004**
- ◆ Monti et al, 1999
- ◆ Chafetz et al, 1962
- ◆ Dauer et al, 2006


# Limitations of published studies

- High refusal rates
- Low statistical power
- Control groups receiving more attention than standard care
- Most studies found positive outcome to some extent in the control group

# Study objectives


1. To test the efficacy of BAI for patients admitted to the ED in modifying hazardous drinkers drinking pattern.
2. To test whether the often-found parallel reduction of alcohol use in control groups is due to the effect of assessing alcohol use and related problems acting like a minimal intervention.

# Study Design



# Research assistants

- 7 baseline research assistants (6 master-level psychologists and 1 ED nurse) conducted screening, assessment and BAI.
- Training included a 2-day workshop on motivational interviewing and a 7-day BAI and research procedures training.
- 3 different follow-up research assistants conducted follow-up telephone interviews.





# Screening

- Cholesterol level
- Primary care physician
- Tobacco use
- Drug use
- Depression
- Immunization
- **Alcohol**
  - ◆ Quantity
  - ◆ Frequency
  - ◆ Frequency of heavy drinking episodes (♂ : > 4 drinks; ♀ [♂ > 65 +] > 3 drinks).

# Assessment (except for control group w/out assessment)

Variables considered for these analyses were:

- Alcohol use questions of the screening
- Socio-demographic information
- AUDIT (score > 12 considered alcohol dependent).

Relevant variables were determined based on prior BAI studies suggesting that certain subgroups were more likely to benefit from BAI (age, gender, medical condition, alcohol dependence).

# Brief Alcohol Intervention

Using an **empathic style** avoiding any confrontation


1. Thank for participation, reassure about confidentiality and assure that any decision about treatment belongs to the patient.
2. **Give feedback** about alcohol use.
3. **Ask patient to comment about feedback.** Ask permission and provide comment regarding the association between alcohol use and risk of injury or other medical conditions.
4. **Ask about the “pros” and “cons” of individual’s alcohol use.**
5. **Ask about importance to change and readiness to change** on 1-10 scale.
6. **Ask what objective patient feels ready** to complete.
7. Depending on patient’s own objective, **affirm patient’s self-efficacy** to achieve his/her objective.
8. **Give a summary document** including patient’s own:
  - ✓ AUDIT score (using data of the intake assessment)
  - ✓ Percentile AUDIT score compared to the general population
  - ✓ Objectives (timeframe, setting of drinking moderation...).

# Follow-up

The variable considered for these analyses were

- Alcohol use screening questions :
  - ◆ Quantity
  - ◆ Frequency
  - ◆ Frequency of heavy drinking episodes (♂ : > 4 drinks; ♀ [♂ > 65+] > 3 drinks)
- AUDIT

# Results - Sample



# Patients' characteristics

- 1366 patients, including 1064 men (78 %) and 302 women (22%)
- Mean age 38.7 (17.31) years
- 68 % Swiss
- 59 % employed
- 987 (72 %) trauma
- 379 (28 %) other surgery (general, urology neurosurgery, other)

## Patients' Characteristics by Group at Intake

	Brief alcohol intervention	Control with assessment	Control w/out assessment	P value
<b>N = 1366</b>	<b>486</b>	<b>543</b>	<b>337</b>	
% Men	76.1	79.0	78.6	0.50
% 18-30	43.4	41.0	51.3	<.01
% Swiss	68.1	68.3	-	0.73
% Employed	59.3	59.1	-	1.00
# Days drinking per week (last 12-mo) (SD)	3.7 (2.4)	3.8 (2.5)	3.6 (2.4)	0.48
# Drinks per drinking occasion (last 12-mo) (SD)	4.3 (3.1)	4.0 (2.7)	3.9 (2.4)	0.24
# Heavy drinking episodes per mo (last 12-mo) (SD)	4.9 (7.4)	4.6 (7.1)	4.1 (6.3)	0.26
AUDIT score (SD)	9.4 (4.7)	8.8 (5.1)	-	0.06

# Alcohol Use Characteristics by Group at Follow-up

	Brief alcohol intervention	Control with assessment	Control w/out assessment	P value
<b>N = 1055 (77.2 %)</b>	<b>367</b>	<b>429</b>	<b>259</b>	
<b>12-month follow-up data</b>				
# Days drinking per week (last 12-mo) (SD)	3.3 (2.3)	3.4 (2.5)	3.1 (2.4)	0.29
# Drinks per drinking occasion (last 12-mo) (SD)	3.5 (2.6)	3.4 (2.5)	3.4 (2.5)	0.63
# Heavy drinking episodes per mo (last 12-mo) (SD)	3.7 (6.0)	3.6 (6.3)	3.6 (6.4)	0.98
AUDIT score (SD)	7.5 (4.7)	7.0 (4.3)	7.3 (4.7)	0.32



# Intake to 12-month Difference in Drinking Pattern

	<b>Brief alcohol intervention</b>	<b>Control with assessment</b>	<b>Control w/out assessment</b>	<b>P value</b>
<b>N = 1055</b>	<b>486</b>	<b>543</b>	<b>337</b>	
# Days drinking per week (last 12-mo) (SD)	-0.4 (1.8)	-0.4 (1.8)	-0.5 (2.0)	0.59
# Drinks per drinking occasion (last 12-mo) (SD)	-0.4 (2.5)	-0.5 (2.8)	-0.4 (2.7)	0.90
# Heavy drinking episodes per mo (last 12-mo) (SD)	-0.7 (7.0)	-0.7 (6.2)	-0.3 (6.8)	0.58
AUDIT score (SD)	-1.8 (3.8)	-1.9 (4.6)	-	0.94
<b>% Changed to low-risk drinking at follow-up</b>	<b>35.7</b>	<b>35.2</b>	<b>37.1</b>	<b>0.88</b>

# 12-month follow-up in subgroups

% changed to low-risk drinking at follow-up

<b>N = 1055</b>		<b>Brief alcohol intervention</b>	<b>Control with assessment</b>	<b>Control w/out assessment</b>	<b>P value</b>
<b>% changed to low-risk drinking at follow-up</b>	<b>N</b>	<b>367</b>	<b>429</b>	<b>259</b>	
	Men	31.6	33.0	33.1	0.92
	18-30 years	35.6	29.7	32.8	0.55
	31-50 years	30.6	33.6	36.9	0.63
	51-65 years	39.7	44.0	50.0	0.63
	66+	45.9	43.2	42.9	0.96
	AUDIT > 12	46.3	37.3	-	0.38
	Trauma	35.6	37.0	37.0	0.71

# GEE models predicting change to low risk drinking at 12 month follow-up

<b>N = 796</b>	<b>Odds-ratio</b>	<b>CI 95 %</b>	<b>Wald</b>	<b>P value</b>
BAI	1.00	[0.74 – 1.33]	0.03	0.87
Men	0.56	[0.41 – 0.76]	14.18	< 0.001
18-30 years	0.96	[0.79 – 1.15]	0.22	0.64
51-65 years	1.47	[1.17 – 1.85]	10.72	0.001
66+	1.57	[1.06 – 2.35]	4.99	0.025
AUDIT > 12	1.54	[1.16 – 2.03]	9.24	< 0.01
Trauma	0.96	[0.74 – 1.24]	0.10	0.76
<b>(Intercept)</b>	0.74	[0.59 – 0.93]	6.69	0.01

- Covariates determined based on prior BAI
- GEE model adjusted for clustering of patients by intake research assistant

# Discussion - Efficacy

## **BAI did not influence a change to low-risk drinking over the 12-month follow-up**

- This null finding applied also for patients previously considered likely to benefit from BAI, i.e., non alcohol-dependent hazardous drinkers and young patients attending the ED after a trauma.
- Limitations to the efficacy of BAI observed may be explained by
  - ◆ The setting: a busy environment, noisy, frequent interruptions may hinder the empathic style of BAI
  - ◆ A large proportion of young patients with minor trauma who may be using ED as a primary care
  - ◆ A single intervention without booster session.

# Discussion – Parallel reduction in control group

**35 % initially hazardous drinkers changed to low-risk drinking at follow-up, also in control groups**

- Two possible explanations for this finding:
  - ◆ a regression to the mean effect
    - ◆ May explain the reduction in alcohol use observed across all groups, but not the absence of an additional effect of BAI on drinking outcomes
  - ◆ an assessment effect
    - ◆ But the study suggests that the alcohol assessment at baseline did not influence drinking pattern over the 12-month follow-up (no difference in outcome between control groups)
    - ◆ A possible explanation may be that already very short, simple screening questions or expectation of follow-up were as efficient as more intense assessment

# Conclusion

- BAI in ED did not influence hazardous drinking
- The reduction of alcohol use in control groups is not due to the effect of assessing alcohol use and related problems
- The positive outcome observed also in control groups may be explained either by a regression to mean effect and/or an effect related to the very short screening or to the expectation of the 12-month follow-up

# Conclusion

- ED may be not be an appropriate setting for BAI
- ED may be more appropriate for screening and referral rather than for BAI itself, as suggested in earlier studies (Chafetz et al, 1962; Crawford et al, 2004)

Thank you!