



UNIVERSITY OF GOTHENBURG

Coached implementation of EIBI


Per Blanck

Fredrik Spak

INEBRIA sept. 2010

The Sahlgrenska Academy

Aim



Implement methods and tools for screening and brief intervention on alcohol problems in the primary health care organisation of the VG-region



Coaches – as implementation **facilitators**

Personel recruited on these qualities:

- Experienced in early intervention
- Trained in MI
- Long practical experience within their profession
- Interested in development work



Evaluation - methods

Process evaluation based on:

Interviews


Documents


Patient surveys



Implementation process

Coach initiatives and support:





Innovation process in an organization, Rogers 2003

Agenda setting

Matching

Redefining/
restructuring

Claryfying

Routinizing



Some numbers – participation in the process

48 **primary health care units**

650 personel , 100 physicians

Maternal health care

55 units 180 nurses

Child health care units

42 units about 200 nurses



Results – typical answers

We have changed our routines, - *maternal health care*

AUDIT has given me a good instrument for further discussions with the pregnant woman – *maternal health care*

Today we discuss alcohol issues with all parents – *child health care*

I am more confident in asking questions about alcohol, as I know how and why I should ask – *child health care*



Results – typical answers

Yes – we discuss alcohol issues among other life style habits - *district nurse*

Some doctors use AUDIT as an instrument with some patients, but I think we should use it much more often – *nurse at general practitioners office*



Results – maternal and child health care

- Alcohol accepted as an important issue
- Confidence in methodology
- Very few patients at risk; one or two per year



Results - numbers

Maternal health care – all pregnant women take an full AUDIT-test

Child health care – all parents are briefly informed and given the opportunity to discuss alcohol issues concerning the relation parent/child



Results – patient survey

“Have your doctor discussed alcohol habits during this consultation?”

Proportion of patients answer “Yes”	Proportion of primary health care units
1-4	14
5-9	45
10-14	24
15-	17

Median – 9 %, n=3 000

Difficult to define risky drinking

Almost none of the interviewed
refers to official guidelines

A simple and well known
definition is needed

Summary

Routines has been changed in a disired direction

Staff feel more secure in addressing these questions with patients.

Systematic screening is used in a limited extent

Risky drinking definition is unclear

Coach – is a good idea when implementing