

**Identifying Core Behavioral and
Psychosocial Data Elements relevant to
substance use disorders for the
Electronic Health Record (EHR)**

Udi Ghitza, Ph.D.

Health Scientist Administrator

Center for the Clinical Trials Network

NIDA/NIH

8th Annual Conference of INEBRIA

September 22, 2011

Immediate CTN EHR Project Goals

- Incorporate consensus-based common data elements relevant to screening and treatment of substance use disorders into a public data standards repository database as a resource for biomedical researchers and EHR vendors.

Consider:

- Office of the National Coordinator for Health Information Technology (ONC) **common data standards** requirements
- CMS requirements for “**meaningful use**”
- Code selected common data elements in terms of established vocabularies and data standards consistent with those of ONC and CMS
- Deposit these common data elements into the NCI’s Cancer Data Standards Registry and Repository (caDSR)
- Our current effort of identifying consensus-based common data elements relevant to screening and treatment of substance use disorders has been informed by 7 meetings at national conferences with federal, state, professional society and community-based treatment stakeholders. An example follows.

**Summary: May 2-3 NIH OBSSR
and Society of Behavioral Medicine
Meeting on “Identifying Core
Behavioral and Psychosocial Data
Elements for the Electronic Health
Record”**

Background

- This NIH meeting was organized and supported by the Office of Behavioral and Social Sciences Research (OBSSR) and Society of Behavioral Medicine.
- Goal: To develop a consensus among various federal, state, community and industry stakeholders on:
 - Recommendations for standardized, practical set of behavioral and psychosocial patient report variables to be collected in primary care electronic health record systems (EHRs)
- Could lead to advancements in data harmonization in behavioral health research.

Background

- **Harmonized screening and collection of standard data on behavioral issues will facilitate:**
 - **Screening, brief interventions and appropriate referrals to specialty treatment in primary care**
 - **Patient-centered shared clinical decision-making**
 - **Population health management**
 - **Biomedical research**
 - **Comparative Effectiveness Research**
 - **Epidemiology**
 - **Personalized medicine (through meta-analyses of large data sets combining health behavior data with medical and biological information)**

Substance Use Disorders: General Recommendations

- **Illicit drug, alcohol and tobacco use should be screened (i.e., screening, brief intervention and referral to treatment) in the primary care setting.**
- **Employ a staged approach to screening and assessment**
 - **Employ single-question screeners for illicit drug and alcohol use (validated where available)**
 - **Employ additional brief assessment instruments (validated where available)**
- **Future considerations**
 - **Employ additional substance use disorder screening assessment questions from standardized instruments where there is significant alcohol or illicit substance use**
 - **Employ longitudinal data collection with standardized questions and timeframe (for example, 3 months)**

Substance Use Disorders: Specific Recommendations

- **Specific Recommendations for core validated drug use screening questions for EHR:**
 - **Single-question screening for illicit drug use and prescription medication use for non-medical reasons:**
 - **Smith PC, Schmidt SM, Allensworth-Davies D, and Saitz R. A single-question screening test for drug use in primary care. *Arch Intern Med* 2010; 170(13):1155-1160**
“How many times in the past year have you used an illegal drug or used a prescription medication for non-medical reasons?”
 - **For additional screening/assessment, the DAST-10 was chosen over the NM-ASSIST because of brevity and the extent of validation:**
 - **Yudko E, Lozhkina O, Fouts A. A comprehensive review of the psychometric properties of the Drug Abuse Screening Test.**
J Subst Abuse Treat 2007;32(2):189-198

Risky Drinking/Alcohol Use: Specific Recommendations

- **Specific Recommendations for core validated risky drinking/ alcohol use disorder questions for EHR:**
 - **Pre-screen question: “Do you sometimes drink alcoholic beverages?”**
 - **Single Question Alcohol Use Screener:**
 - **Smith PC, Schmidt SM, Allensworth-Davies D and Saitz R. Primary care validation of a single-question alcohol screening test. *J Gen Intern Med* 2009; 24(7):783-8 (NIAAA recommended question)**
- **“How many times in the past year have you had X or more drinks in a day? (where X is 5 for men and 4 for women, and a response of ≥ 1 is considered positive).”**
 - **Additional assessment: Alcohol Use Disorders Identification Test – Consumption (3-item AUDIT-C):**
 - **Bush K, Kivlahan DR, McDonell MB, Fihn SD and Bradley KE. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol use disorders identification test. *Arch Intern Med* 1998; 158:1789-95**

Tobacco Use: Specific Recommendations

- **Specific Recommendations for core validated tobacco use questions for EHR:**
 - **Implement screening and assessment with validated questions for the 5 A's.**
 - **Additional suggestions:**
 - **Use BRFSS – Behavioral Risk Surveillance System (CDC) for the “Ask” question regarding identifying and documenting tobacco use status for every patient**
 - **Ask about exposure to secondary smoke**
 - **Other screening instruments commented in the wiki, such as the FTND, were not considered to be appropriate for primary care.**

Tobacco Use: Specific Recommendations

- Recommended Screener questions for all patients at every visit

Have you used tobacco in the last 30 days?

Smoked Cigarettes: Yes ____ No ____

Used a Smokeless Tobacco Product:

Yes ____ No ____

For all who responded YES to having smoked or used smokeless tobacco in the last 30 days:

- Would you be interested in quitting tobacco use within the next few weeks?


Yes ____ No ____


For all who responded YES to Question 2, the EHR then prompts the CLINICIAN to answer the following questions:

Did you provide brief counseling/coaching to quit? Yes ____ No ____

Did you prescribe or recommend that the patient to use one of the seven FDA-approved medications for tobacco cessation? Yes ____ No ____

Did you refer the patient to your State's Tobacco Quitline (1-800-QUIT-NOW):
Yes ____ No ____





EMR Workgroup: Acknowledgements



■ EMMES Corp.

- Bob Lindblad
- Robert Gore-Langton
- David Shide
- Paul Van Veldhuisen
- Brian Campbell

■ NIDA CTN


- John Rotrosen
- CTN Special Interest
Group on EMR
- CTN Community Treatment Programs (CTPs)

■ NIDA CCTN

- Betty Tai
- Steve Sparenborg
- Geetha Subramaniam
- Paul Wakim
- Carol Cushing

■ NIDA DESPR

- Sarah Duffy



CTN EMR SIG Members

- **John Rotrosen**
- **Carla Green**
- **Jennifer McNeely**
- **Andy Saxon**
- **Dennis McCarty**
- **Ron Jackson**
- **Daniel Kivlahan**
- **Marc Gourevitch**
- **Harold Pincus**
- **Ken Gersing**
- **John March**
- **Denny Carise**
- **Dave Oslin**
- **Al Cohen**
- **Larry Brown**
- **Madhukar Trivedi**
- **George Woody**